arXiv:1304.6989v1 [cond-mat.quant-gas] 25 Apr 2013

Tests of universal three-body physics in an ultracold Bose-Fermi mixture

Ruth S. Bloom, Ming-Guang Hu, Tyler D. Cumby, and Deborah S. Jin
JILA, NIST and University of Colorado; Department of Physics, University of Colorado, Boulder
(Dated: April 26, 2013)

Recent measurements of Efimov resonances in a number of ultracold atom species have revealed an unexpected universality, in which three-body scattering properties are determined by the van der Waals length of the two-body interaction potential. To investigate whether this universality extends to heteronuclear mixtures, we measure loss rate coefficients in an ultracold trapped gas of ^{40}K and ^{87}Rb atoms. We find an Efimov-like resonance in the rate of inelastic collisions between ^{40}K^{87}Rb Feshbach molecules and ^{87}Rb atoms. However, we do not observe any Efimov-related resonances in the rates of inelastic collisions between three atoms. These observations are compared to previous measurements by the LENS group of Efimov resonances in a ^{41}K and ^{87}Rb mixture as well as to recent predictions.

Magnetically tunable Feshbach resonances in ultracold atomic gases make it possible to explore a vast array of phenomena [1], and recent years have witnessed a rapid evolution in the understanding of universal few-body effects that accompany resonantly enhanced two-body interactions. First proposed in the context of nuclear theory, the Efimov effect consists of an infinite series of three-body bound states when the two-body scattering length a diverges [2–5]. These Efimov trimers exist even when the two-body potential does not support a bound state, and their spatial extent can reach far beyond the length scale of inter-particle forces. Most notably, Efimov trimers follow a discrete scale invariance, with each subsequent bound state larger than the last by a scaling factor of $e^{\pi/s_0} \approx 22.7$ for identical bosons.

Several ultracold atom experiments have now leveraged tunable interactions to measure Efimov physics [6–15]. In these dilute gases, weakly bound Efimov trimer states mediate inelastic collisions and give rise to a rich spectrum of atom loss resonances as a function of a [4]. A surprising result of these experiments has been the emergence of a pattern in the locations of the Efimov loss resonances for homonuclear trimers, which consistently appear at $a \approx -9.1\, l_{\mathrm{vdW}}$ for different nuclear spin states [13], different Feshbach resonances [14], and different atomic species [7–9, 11, 13–15], where l_{vdW} is the two-body van der Waals length. The three-body parameter that sets the locations of Efimov resonances was previously understood to be a non-universal quantity subject to the unique details of each system's interaction potential at short range [3, 4, 16]. However, recent theoretical work has shown that van der Waals potentials in cold atoms are responsible for screening these non-universal details [17–21]. A universal three-body parameter has also been predicted for heteronuclear mixtures [22].

Extending the study of Efimov trimers to mixtures of atom species is of particular interest in exploring universal three-body physics [1, 10, 23–25]. In these systems, only two of the three pairwise interactions are resonant and the Efimov scaling factor depends on the mass ratio of the constituent particles [4]. Halo nuclei, in which two neutrons orbit a heavier nuclear core at a distance well beyond the classically allowed radius, are an example of Efimov-like three-body physics for a mass-imbalanced system [26, 27]. However, in cold atom systems there has been only one clear measurement of mixed-species Efimov trimers to date, in work carried out by the LENS group for a mixture of ^{41}K and ^{87}Rb atoms [10].

In this Letter, we investigate Efimov physics in a second quantum gas mixture, namely ^{40}K and ^{87}Rb atoms, where we measure rates of inelastic loss near an interspecies Feshbach resonance. We report the observation of an Efimov-like resonance in collisions between ^{87}Rb atoms and ^{40}K^{87}Rb Feshbach molecules, but no observed Efimov resonances in three-body atom loss. If the three-body parameter is universal for heteronuclear trimers, then Efimov resonances connected to RbRbK trimers should occur at similar values of a for mixtures of ^{40}K ad ^{87}Rb and for mixtures of ^{41}K ad ^{87}Rb [22]. Therefore, we compare our measurements with the previous observations by the LENS group [10].

Our measurements of inelastic loss start with an ultracold gas mixture of bosonic ^{87}Rb atoms in the $|f, m_f\rangle = |1, 1\rangle$ state and fermionic ^{40}K atoms in the $|9/2, -9/2\rangle$ state, which are the lowest energy hyperfine states in a magnetic field. Here, f corresponds to the total atomic angular momentum and m_f is its projection. The atom gas is prepared near the onset of quantum degeneracy, with the temperature T above 1.1 T_{c} for ^{87}Rb and above 0.2 T_{F} for ^{40}K, where T_{c} is the transition temperature for Bose-Einstein condensation and T_{F} is the Fermi temperature. The gas is confined in a far-detuned optical dipole trap formed by a focused beam of 1090 nm light. The radial and axial trap frequencies for ^{87}Rb range from 400 to 500 Hz and 6 to 10 Hz, respectively, with corresponding frequencies for ^{40}K higher by a factor of 1.4. By measuring the resonant absorption of probe beams directed along a radial trap direction, both atom species are imaged within a few milliseconds of their release from the optical trap. In these images, the atom distribution is nearly in-situ along the weak axial trap direction, and fully in momentum-space along the tight radial direc-



FIG. 1. (Color online) β vs. a, for collisions of ^{40}K^{87}Rb molecules with ^{40}K atoms in the $|9/2, -7/2\rangle$ state (open triangles), with ^{40}K atoms in the $|9/2, -9/2\rangle$ state (open squares), and with ^{87}Rb atoms in the $|1, 1\rangle$ state (solid diamonds). Vertical error bars correspond to one standard deviation and represent statistical error in β. Horizontal error bars correspond to a ± 0.02 G error in B, which is an upper bound on the variation in B across the cloud. A broad resonance is evident in the ^{87}Rb+^{40}K^{87}Rb loss rate data. The solid curve shows Eq. (2) with Efimov parameters $a_* = 230(10)\ a_0$, $\eta_* = 0.26(3)$, and $C_\beta = 3.2(2)$.

tion. In-trap density profiles are determined from fits to Maxwell-Boltzmann distributions, assuming ballistic expansion. An s-wave magnetic Feshbach resonance is used to control interactions between the ^{87}Rb atoms and the ^{40}K atoms, where the scattering length at a magnetic field B near the resonance is given by $a = a_{\rm bg}(1 - \frac{\Delta}{B - B_0})$, with $a_{\rm bg} = -187\ a_0$, $B_0 = 546.618$ G, $\Delta = -3.04$ G, and a_0 is the Bohr radius [28].

We begin by considering inelastic atom-molecule collisions. Weakly bound molecules, which exist on the positive a side of the Feshbach resonance, can undergo collisional vibrational relaxation into deeply bound states. The energy released in this process is sufficiently large that both collision partners are lost from the trap. The inelastic collision rate is enhanced when a is tuned to a value where an Efimov trimer state is degenerate with the atom-molecule threshold [4, 29]. Such Efimov resonances have been observed as broad loss features for ^{133}Cs [30] and ^{6}Li dimers [12, 31].

For the case of ^{87}Rb atoms and ^{40}K^{87}Rb Feshbach molecules, Helfrich *et al.* have suggested that an Efimov resonance exists just outside the range of previous loss measurements [24, 32]. To determine whether an Efimov resonance is in fact present, we measure molecule loss over a wider range of a. We compare results for molecules colliding with three different partners: ^{87}Rb atoms in the $|1, 1\rangle$ state, ^{40}K atoms in the $|9/2, -9/2\rangle$ state, and ^{40}K atoms in the $|9/2, -7/2\rangle$ state. Here, Efimov resonances are only possible for the first case, and, in the last case, the atom collision partner does not participate in the Feshbach resonance.

For these measurements, the atom gas mixture is typically prepared at $a = -300\ a_0$ and $T = 300$ nK. The magnetic field is slowly swept across the Feshbach resonance at a rate of 3 G/ms to form KRb Feshbach molecules by magnetoassociation [33]. In order to prepare a pure sample of molecules and one particular atom species, we use a combination of resonant light and RF pulses to selectively remove unpaired ^{40}K, or ^{87}Rb, atoms from the trap. The ^{87}Rb atom removal is most efficient for a small sample, so K+KRb collisions are measured with an additional confining laser beam that increases the K axial trap frequency to 30 Hz. For the case of non-resonant atom-molecule collisions, we transfer the ^{40}K atoms into the $|9/2, -7/2\rangle$ hyperfine state using RF adiabatic rapid passage.

To measure trap loss rates, we sweep the magnetic field to the value desired for the loss measurement at a rate that is fast in comparison to loss timescales, wait a variable hold time, turn off the optical trap, and then jump the magnetic field back across the Feshbach resonance at a rate of 50 G/ms to dissociate the molecules. From images of the resulting atom clouds, we extract the number of molecules, the number of free atoms, and the in-trap cloud sizes [34]. The atom-molecule collisions are characterized by a loss rate coefficient β, which is defined by

$$\dot{N}_{\rm KRb}(t) = \dot{N}_{\rm A}(t) = -\beta \int d^3\mathbf{r}\ n_{\rm KRb}(\mathbf{r}, t)\ n_{\rm A}(\mathbf{r}, t), \quad (1)$$

where $n_{\rm KRb}(\mathbf{r}, t)$ is the number density of the molecules and $n_{\rm A}(\mathbf{r}, t)$ is the number density of the atoms. We extract β by fitting the measured loss of atoms and molecules to numerical solutions of coupled Eqs. (1). These solutions account for heating by including the measured cloud sizes at each point in time. We estimate that the systematic uncertainty in β, which is limited by the uncertainty in the measured in-trap densities, is $\pm 40\%$.

The plot of β vs. a in Fig. 1 reveals a resonance in the collisions between KRb molecules and Rb atoms, which we attribute to an Efimov-like RbRbK bound state. This is consistent with the absence of a resonance in the loss of KRb molecules to collisions with K atoms in either hyperfine state. The universal shape of an atom-molecule Efimov resonance at $T = 0$ is given by Ref. [24],

$$\beta(a) = C_\beta \frac{\sinh(2\eta_*)}{\sin^2[s_0\ \ln(a/a_*)] + \sinh^2(\eta_*)} \frac{\hbar a}{m_1}. \quad (2)$$

Here, η_* sets the Efimov resonance width and a_* sets the scattering length at the peak of the Efimov resonance. In the ^{40}K-^{87}Rb mixture, s_0=0.6536 and m_1 is the mass of ^{40}K [24]. The solid line in Fig. 1 shows a fit of



FIG. 2. (Color online) α vs B. There are two clear peaks near 546.6 G and 547.4 G: The first is the s-wave Feshbach resonance that we use to control a, and the second, much narrower feature we attribute to a d-wave interspecies Feshbach resonance. Vertical error bars show the statistical uncertainty in α, and horizontal error bars show an upper bound of ± 0.03 G for the variation in B during the measurement. Inset: α vs. B near the feature at 547.4 G. The solid, dotted, and dashed lines show the universal shape of an Efimov resonance, Eqs. (4) and (5), for $\eta* = 0.1$, $\eta* = 0.02$, and $\eta* = 0.004$, respectively. The data are not consistent with the shape of an Efimov resonance for any value of $\eta*$.

Eq. 2 to the measured Rb+KRb loss rate coefficients for $a \geq 140\ a_0$, which gives $a_* = 230(10)\ a_0$, $\eta* = 0.26(3)$, and $C_\beta = 3.2(2)$. The location of the loss resonance is consistent with the suggestion of $a_* = 200(50)\ a_0$ made by Ref. [24]. This reference predicts $C_\beta = 10.1$ for the ^{40}K-^{87}Rb mixture, however we note that a similar difference between theoretical and measured amplitudes has also been seen for single-species atom-molecule loss resonances [12, 30, 31].

We now turn our attention to three-body recombination rates, which are widely used for measuring Efimov physics in cold atoms. Three-body recombination is an inelastic process, in which three atoms collide and form a diatomic molecule, with the remaining atom required for conservation of energy. Since each collision event typically results in the loss of all three atoms from the trap, the inelastic collision rate can be extracted from measurements of atom loss. Three-body recombination is resonantly enhanced when an Efimov trimer state approaches zero binding energy at a negative a denoted a_- [4, 35]. On the other side of the Feshbach resonance, interference effects related to the Efimov states lead to minima in the three-body recombination rate at positive a denoted a_+ [4, 35]. These features modulate an overall a^4 dependence of the three-body recombination rate that results in a large increase of atom loss rates near a Feshbach resonance [4, 35].

In order to avoid crossing the Feshbach resonance, we prepare the atom gas mixture by evaporating at a field where $|a| \approx 300\ a_0$ on the same side of the Feshbach resonance as the field for the loss measurement, B. The magnetic field is then swept to B in a time that is short compared to the loss, and held for a variable time t. Finally, the optical trap is turned off and the field is swept back to the evaporation value where the atoms are imaged.

Two collision channels exist for interspecies three-body recombination, which in our case are ^{87}Rb+^{87}Rb+^{40}K and ^{87}Rb+^{40}K+^{40}K. Only the first one supports Efimov resonances [4], and the latter one involving two identical fermions is suppressed by Fermi statistics. For the first channel, an event rate coefficient α is defined by,

$$\dot{N}_{\mathrm{Rb}}(t) = 2\dot{N}_{\mathrm{K}}(t) = -2\alpha \int d^3\mathbf{r}\ n_{\mathrm{K}}(\mathbf{r},t)\ n_{\mathrm{Rb}}^2(\mathbf{r},t). \quad (3)$$

Here, $n_{\mathrm{Rb}}(\mathbf{r},t)$ and $n_{\mathrm{K}}(\mathbf{r},t)$ are the densities of ^{87}Rb and ^{40}K, respectively. From our measured cloud sizes and atom numbers as a function of t, we obtain α using an analysis similar to that described in Ref. [36], which uses the measured atom densities at each time in order to account for heating of the gas. We estimate the systematic uncertainty in α, which dominated by the uncertainty for measured in-trap densities, to be $\pm 20\%$.

In Fig. 2, we show the measured α as a function of B. We measure an increase of α by more than four orders of magnitude at the s-wave resonance at B_0=546.618 G. A second peak, on the order of 0.1 G wide, is also evident near 547.4 G. As illustrated in the inset to Fig. 2, this feature is not consistent with the expected shape of an Efimov resonance [24], and we identify it as a two-body d-wave Feshbach resonance [37, 38]. Because it is technically challenging to control a near a narrow resonance, we exclude the data for B between 547.25 G and 547.50 G from further analysis.

The lower panel of Fig. 3 shows the measured α vs. a. For comparison, we also plot the universal form for α as a function of a [24] with Efimov resonances at hypothetical locations. At negative a,

$$\alpha(a<0) = \frac{C_\alpha}{2} \frac{\coth(\pi s_0)\sinh(2\eta_*)}{\sin^2[s_0\ \ln(a/a_-)] + \sinh^2(\eta_*)} \frac{\hbar a^4}{m_1}, \quad (4)$$

whereas for positive a,

$$\alpha(a>0) = C_\alpha \left(\frac{\sin^2[s_0\ \ln(a/a_+)] + \sinh^2\eta_*}{\sinh^2(\pi s_0 + \eta_*) + \cos^2[s_0\ \ln(a/a_+)]} \right. \\ \left. + \frac{\coth(\pi s_0)\cosh(\eta_*)\sinh(\eta_*)}{\sinh^2(\pi s_0 + \eta_*) + \cos^2[s_0\ \ln(a/a_+)]} \right) \frac{\hbar a^4}{m_1}, \quad (5)$$

where the coefficient C_α is 345 [24]. For the curves in Fig. 3, we use $\eta* = 0.02$, which gives the best agreement



FIG. 3. (Color online) Lower panel: α vs. a. The error bars correspond to those in Fig. 2. The solid line shows an a^4 dependence, which is expected in the absence of Efimov resonances. In order to clarify the range of a for which Efimov features are excluded, we include dashed and dash-dotted lines corresponding to universal Efimov resonance shapes located at $a_- = -200\ a_0$ and $a_- = -3000\ a_0$, respectively. Similarly, the dash-dot-dotted and dotted lines show hypothetical Efimov interference minima at $a_+ = 3000\ a_0$ and $a_+ = 200\ a_0$, respectively. We conclude that there are no Efimov features in the ^{40}K-^{87}Rb mixture for $-3000 \leq a_-/a_0 \leq -200$ and $200 \leq a_+/a_0 \leq 3000$. A narrow two-body Feshbach resonance blocks experimental access to the range $-1090 \leq a/a_0 \leq -840$ (shaded rectangles on axes). Upper panel: The ratio of Rb loss to K loss vs. a. The solid line shows $\Delta N_{\mathrm{Rb}}/\Delta N_{\mathrm{K}} = 2$, which is expected for three-body recombination involving one ^{40}K atom and two ^{87}Rb atoms.

to the overall magnitude of α in the measurements. We conclude that there are no Efimov features at negative a between -200 a_0 and -3000 a_0 or at positive a between 200 a_0 and 3000 a_0.

The upper panel of Fig. 3 shows the measured ratio of ^{87}Rb loss to ^{40}K loss. Within the uncertainties, this ratio does not show a dependence on a, and the average value is 2.1(1). This is consistent with the dominant loss process involving two ^{87}Rb atoms and one ^{40}K atom.

While the data in Fig. 3 cover a factor of twenty in a, this range is small compared to the Efimov scaling factor of $e^{\pi s_0} = 122.7$ for ^{40}K-^{87}Rb-^{87}Rb [24], and therefore it is perhaps not surprising that we do not observe any Efimov features. It is experimentally challenging to extend the range of a where universal Efimov physics can be explored. On the low $|a|$ side, this range is limited by the requirement that $|a|$ be large compared to the physical range of the interactions, which can be characterized by the magnitude of the effective range R_{e} [39–41], where $R_{\mathrm{e}} = -140\ a_0$ for ^{40}K-^{87}Rb [1]. For higher $|a|$, the useful range of a is limited to $k|a| < 1$, where k is the thermal wave number, $k \approx \sqrt{2\mu_{\mathrm{KRb}} k_B T}/\hbar$, k_{B} is the Boltzmann constant, and μ_{KRb} is the reduced mass for ^{40}K and ^{87}Rb [42]. Our measurements of α at high $|a|$ are typically near $T = 300$ nK, which gives $1/k_{\mathrm{th}} = 3,000\ a_0$.

One motivation for our work was the fact that heteronuclear Efimov resonances have previously been seen in a mixture of ^{41}K and ^{87}Rb atoms, where the LENS group found a peak in α at a_-=$-249\ a_0$ due to an ^{87}Rb+^{87}Rb+^{41}K resonance [10]. Given the similarity of ^{87}Rb+^{87}Rb+^{41}K and ^{87}Rb+^{87}Rb+^{40}K in mass ratios and van der Waal lengths, one would expect similar locations for Efimov resonances in these two systems if the three-body parameter is universally determined by the two-body potential; however, we do not observe a resonance in our data (Fig. 3). Given the relatively small $|a|$ for the observed ^{87}Rb+^{87}Rb+^{41}K resonance, it is possible that finite-range effects could play a role in explaining the absence of a resonance near this a_- for ^{87}Rb+^{87}Rb+^{40}K.

Finally, we can compare our measurements to universal predictions for Efimov resonance locations in heteronuclear mixtures. Helfrich *et al.* in Ref. [24] predict $a*/|a_-| = 0.51$ for the ^{40}K and ^{87}Rb system. Based on our observed atom-molecule resonance, this would put an Efimov resonance in α at $a_- = -430\ a_0$, which is inconsistent with our measurements of three-body loss. However, we note that even for three identical bosons, measurements of $a*/|a_-|$ have so far not agreed with universal Efimov theory [12, 30, 31].

Wang *et al.* have generalized the three-body parameter's universality to heteronuclear mixtures and predict the absolute positions of Efimov resonances for three-body loss in K+Rb mixtures to be $a_+ = 2800\ a_0$ and $a_- < -30,000\ a_0$ [22]. These values are at or beyond the limits of our experimental range, which is consistent with the absence of loss features in our measurements of α.

In conclusion, with an ultracold gas mixture of ^{40}K and ^{87}Rb atoms, we observe an Efimov-like resonance in atom-molecule inelastic collisions, but no Efimov feature in three-body recombination. In comparing this with previous data for a mixture of ^{41}K and ^{87}Rb atoms, we do not find evidence for universality of the three-body parameter. Nevertheless, our data is consistent with a recent prediction, which is based on a universal three-body parameter, that puts the first universal Efimov resonance outside the reach of our experiments at $a_- < -30,000\, a_0$. Additional data, including measurements for different systems, will help to address the question of universality in the locations of Efimov resonances for heteronuclear atom gas mixtures. In particular, heteronuclear systems with smaller scaling parameters offer the best chance for observing multiple Efimov resonances within an experimentally accessible range of $|a|$. Another direction for future work is to RF-associate atoms into a heteronuclear trimer state, in order to map out the trimer binding energy as a function of a [43].

The authors thank Chris Greene, Jose D'Incao, John Bohn, Yujun Wang, and Eric Cornell for discussions. RSB acknowledges support from NSF-GRFP and NDSEG. This work was funded by NIST and the NSF.

[1] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Rev. Mod. Phys. **82**, 1225 (2010).

[2] V. N. Efimov, Yad. Fiz. **12**, 1080 (1970).

[3] V. Efimov, Nuclear Physics A **210**, 157 (1973).

[4] E. Braaten and H.-W. Hammer, Phys. Rep. **428**, 259 (2006).

[5] N. T. Zinner and A. S. Jensen, J. Phys. G **40**, 053101 (2013).

[6] T. Kraemer, M. Mark, P. Waldburger, J. G. Danzl, C. Chin, B. Engeser, A. D. Lange, K. Pilch, A. Jaakkola, H.-C. Nägerl, and R. Grimm, Nature **440**, 315 (2006).

[7] T. B. Ottenstein, T. Lompe, M. Kohnen, A. N. Wenz, and S. Jochim, Phys. Rev. Lett. **101**, 203202 (2008).

[8] J. H. Huckans, J. R. Williams, E. L. Hazlett, R. W. Stites, and K. M. O'Hara, Phys. Rev. Lett. **102**, 165302 (2009).

[9] M. Zaccanti, B. Deissler, C. D'Errico, M. Fattori, M. Jona-Lasinio, S. Müller, G. Roati, M. Inguscio, and G. Modugno, Nature Physics **5**, 586 (2009).

[10] G. Barontini, C. Weber, F. Rabatti, J. Catani, G. Thalhammer, M. Inguscio, and F. Minardi, Phys. Rev. Lett. **103**, 043201 (2009).

[11] S. E. Pollack, D. Dries, and R. G. Hulet, Science **326**, 1683 (2009).

[12] S. Nakajima, M. Horikoshi, T. Mukaiyama, P. Naidon, and M. Ueda, Phys. Rev. Lett. **105**, 023201 (2010).

[13] N. Gross, Z. Shotan, S. Kokkelmans, and L. Khaykovich, Phys. Rev. Lett. **105**, 103203 (2010).

[14] M. Berninger, A. Zenesini, B. Huang, W. Harm, H.-C. Nägerl, F. Ferlaino, R. Grimm, P. S. Julienne, and J. M. Hutson, Phys. Rev. Lett. **107**, 120401 (2011).

[15] R. J. Wild, P. Makotyn, J. M. Pino, E. A. Cornell, and D. S. Jin, Phys. Rev. Lett. **108**, 145305 (2012).

[16] J. P. D'Incao, C. H. Chreene, and B. Esry, J. Phys. B **42**, 044016 (2009).

[17] C. Chin, arXiv:1111.1484v2 [**cond-mat.quant-gas**] (2011).

[18] P. Naidon, S. Endo, and M. Ueda, arXiv:1208.3912v1 [**cond-mat.quant-gas**] (2012).

[19] J. Wang, J. P. D'Incao, B. D. Esry, and C. H. Greene, Phys. Rev. Lett. **108**, 263001 (2012).

[20] R. Schmidt, S. P. Rath, and W. Zwerger, Eur. Phys. B **85** (2012), 10.1140/epjb/e2012-30841-3.

[21] P. K. Sørensen, D. V. Fedorov, A. S. Jensen, and N. T. Zinner, Phys. Rev. A **86**, 052516 (2012).

[22] Y. Wang, J. Wang, J. P. D'Incao, and C. H. Greene, Phys. Rev. Lett. **109**, 243201 (2012).

[23] B. Marcelis, S. J. J. M. F. Kokkelmans, G. V. Shlyapnikov, and D. S. Petrov, Phys. Rev. A **77**, 032707 (2008).

[24] K. Helfrich, H.-W. Hammer, and D. S. Petrov, Phys. Rev. A **81**, 042715 (2010).

[25] Y. Wang, W. B. Laing, J. von Stecher, and B. D. Esry, Phys. Rev. Lett. **108**, 073201 (2012).

[26] A. S. Jensen, K. Riisager, D. V. Fedorov, and E. Garrido, Rev. Mod. Phys. **76**, 215 (2004).

[27] T. Frederico, A. Delfino, L. Tomio, and M. Yamashita, Prog Part Nucl Phys. **67**, 939 (2012).

[28] C. Klempt, T. Henninger, O. Topic, M. Scherer, L. Kattner, E. Tiemann, W. Ertmer, and J. J. Arlt, Phys. Rev. A **78**, 061602 (2008).

[29] E. Nielsen, H. Suno, and B. D. Esry, Phys. Rev. A **66**, 012705 (2002).

[30] S. Knoop, F. Ferlaino, M. Mark, M. Berninger, H. Schöbel, H.-C. Nägerl, and R. Grimm, Nat. Phys. **5**, 227 (2009).

[31] T. Lompe, T. B. Ottenstein, F. Serwane, K. Viering, A. N. Wenz, G. Zürn, and S. Jochim, Phys. Rev. Lett. **105**, 103201 (2010).

[32] J. J. Zirbel, K.-K. Ni, S. Ospelkaus, J. P. D'Incao, C. E. Wieman, J. Ye, and D. S. Jin, Phys. Rev. Lett. **100**, 143201 (2008).

[33] T. D. Cumby, R. A. Shewmon, M.-G. Hu, J. D. Perreault, and D. S. Jin, Phys. Rev. A **87**, 012703 (2013).

[34] The atom and molecule clouds are fit to gaussian distributions, and the in-trap radial sizes are calculated from the measured axial size and the trap aspect ratio.

[35] B. D. Esry, C. H. Greene, and J. P. Burke, Phys. Rev. Lett. **83**, 1751 (1999).

[36] C. Ospelkaus, S. Ospelkaus, K. Sengstock, and K. Bongs, Phys. Rev. Lett. **96**, 020401 (2006).

[37] M. Zaccanti, C. D'Errico, F. Ferlaino, G. Roati, M. Inguscio, and G. Modugno, Phys. Rev. A **74**, 041605 (2006).

[38] B. P. Ruzic, C. H. Greene, and J. L. Bohn, Phys. Rev. A **87**, 032706 (2013).

[39] V. Efimov, Phys. Rev. C **44**, 2303 (1991).

[40] M. Thøgersen, D. V. Fedorov, and A. S. Jensen, Phys. Rev. A **78**, 020501 (2008).

[41] L. Platter, C. Ji, and D. R. Phillips, Phys. Rev. A **79**, 022702 (2009).

[42] J. P. D'Incao, H. Suno, and B. D. Esry, Phys. Rev. Lett. **93**, 123201 (2004).

[43] T. Lompe, T. B. Ottenstein, F. Serwane, A. N. Wenz, G. Zürn, and S. Jochim, Science **330**, 940 (2010).